

Free degrees of homeomorphisms and periodic maps on closed surfaces

Shicheng Wang*

Department of Mathematics, Peking University, Beijing 100871, People's Republic of China

Received 17 December 1990

Revised 29 May 1991

Abstract

Wang, S., Free degrees of homeomorphisms and periodic maps on closed surfaces, *Topology and its Applications* 46 (1992) 81–87.

For each closed hyperbolic (orientable or nonorientable) surface F , we provide a positive integer $h(F)$ with properties that for any homeomorphism f on F , at least one of the iterates $f, f^2, \dots, f^{h(F)}$ has a fixed point and there is a periodic (orientation reversing) homeomorphism s on F such that $s^1, s^2, \dots, s^{h(F)-1}$ are all fixed point free. Those integers $h(F)$ are: $h(F_2) = 4$, $h(F_g) = 2g - 2$, $g > 2$, $h(N_3) = 2$, $h(N_q) = q - 2$, $q > 3$. There is an interesting difference between our result and the Nielsen's result in the orientable category.

Keywords: Fixed point, closed surface, homeomorphism, periodic map.

AMS (MOS) Subj. Class.: Primary 55M20, 57A10.

Let M be a closed manifold and s be a homeomorphism on M . The free degree of s is the maximum positive integer m such that s^1, s^2, \dots, s^{m-1} are all fixed point free. We call the maximum free degree of homeomorphisms on M , denoted by $h(M)$, the free degree of homeomorphisms on M .

By the result in [3], $h(M)$ exists if the Euler number of M is nonzero. If M is a differential manifold, by the well-known fact in differential topology, $h(M)$ exists if and only if the Euler number of M is nonzero.

We can define $p(M)$, the free degree of periodic homeomorphisms (but arbitrary periods) on M ; $h_0(M)$, the free degree of orientation preserving homeomorphisms on M (if M is orientable); $p_0(M)$, the free degree of orientation preserving periodic homeomorphisms (but arbitrary periods) on M (if M is orientable) in the same way.

Correspondence to: Professor Wang, Department of Mathematics, Peking University, Beijing, 100871, People's Republic of China.

* The author is partially supported by NSFC grants.

By the definitions, we have $p_0(M) \leq h_0(M) \leq h(M)$, $p_0(M) \leq p(M) \leq h(M)$. Let F_g be an orientable closed surface of genus g , $g \geq 2$, N_q be a nonorientable closed surface of genus q , $q \geq 3$. The result in this paper is

Theorem 1.

$$h(F_g) = p(F_g) = \begin{cases} 4, & \text{if } g = 2, \\ 2g - 2, & \text{if } g > 2, \end{cases}$$

$$h(N_q) = p(N_q) = \begin{cases} 2, & \text{if } q = 3, \\ q - 2, & \text{if } q > 3. \end{cases}$$

Remark. (1) Nielsen studied this problem in the forties in the orientable category. His result is

$$h_0(F_g) = \begin{cases} 2 \text{ or } 3, & \text{if } g = 2, \\ 2g - 2, & \text{if } g > 2, \end{cases}$$

and

$$p_0(F_g) = \begin{cases} 2, & \text{if } g = 2, \\ g - 1, & \text{if } g > 2. \end{cases}$$

(See [5], reviewed by Fox, that here $h_0(F_2)$ is 2 or 3 is still unknown.)

Thus it is a little bit surprising that the free degrees of homeomorphisms are realized by periodic maps in our theorem.

(2) The closed surfaces not included in Theorem 1 are the 2-sphere, torus, projective plane and Klein bottle. All these cases are well studied.

Now we give some definitions and results which will be used in the proof of Theorem 1.

(a) Let S be a closed hyperbolic orbifold with n singular points of indices v_1, v_2, \dots, v_n . We denote the underlying space of S by $|S|$. The general Euler number of S , $\tilde{\chi}(S)$, is defined by the formula

$$\tilde{\chi}(S) = \chi(|S|) - \sum_{i=1}^n \left(1 - \frac{1}{v_i}\right);$$

here χ is the ordinary Euler number. The general fundamental group (or the fundamental group of orbifolds) of S is

$$\pi_1^0(S) = \begin{cases} \langle a_1, b_1, \dots, a_g, b_g, c_1, \dots, c_n \mid c_1^{v_1} = \dots = c_n^{v_n} \\ \quad \quad \quad = [a_1, b_1] \cdots [a_g, b_g] c_1 \cdots c_n = 1 \rangle, \text{ or} \\ \langle a_1, \dots, a_g, c_1, \dots, c_n \mid c_1^{v_1} = \dots = c_n^{v_n} = a_1^2 \cdots a_g^2 c_1 \cdots c_n = 1 \rangle. \end{cases}$$

If $n = 0$, then $\tilde{\chi}(S) = \chi(S) = \chi(|S|)$ and $\pi_1(|S|) = \pi_1^0(S)$. (See [6, Section 2].)

Any surjective homomorphism $\phi: \pi_1^0(S) \rightarrow Z_d$ determines a cyclic branched cover $p: \tilde{S} \rightarrow S$, here Z_d is a cyclic group of order d and \tilde{S} is a hyperbolic orbifold with $\pi_1^0(\tilde{S}) = \text{kernel } \phi$ (see [1] or [4]). \tilde{S} is a hyperbolic surface if and only if $\text{kernel } \phi$ is a surface group. $\text{kernel } \phi$ is a surface group if and only if c_i and $\phi(c_i)$ have the same order [4, Theorem 3].

(b) For any periodic map f on a closed surface F with $\chi(F) < 0$, there is a hyperbolic metric on F such that f is an isometry under this metric.

Let f be an isometry of order d on a closed hyperbolic surface F with isolated periodic points only, then the quotient space $F/\langle f \rangle$ is a hyperbolic orbifold S and each singular point of index v_i is determined by an isolated periodic point orbit of f with length d/v_i .

Furthermore we have the Riemann–Hurwitz formula (see [4])

$$\frac{\tilde{\chi}(F)}{d} = \tilde{\chi}(S) = \chi(|S|) - \sum_{i=1}^n \left(1 - \frac{1}{v_i}\right).$$

(c) Let $f: F \rightarrow F$ be a self map on a closed surface F . The Lefschetz number $L(f)$ is defined as $\sum_{i=0}^2 (-1)^i \text{trace}(f_{*i})$, here f_{*i} is a homomorphism on $H_i(F, Q)$ induced by f . The Lefschetz Fixed Point Theorem claims that if f is fixed point free, then $L(f)$ is zero (see [2]).

Proof of Theorem 1. The proof of Theorem 1 is elementary. It will be finished by two steps. The first step is to construct a periodic map with the required free degree on each closed surface. The constructions are inspired by our work [9]. The second step is to show the numbers in Theorem 1 are the upper bounds of the free degrees of homeomorphisms on closed surfaces. This step is to apply Fuller's idea (originated from Nielsen, see [2, 3]) to our concrete cases and to make a refinement.

Step 1. There are five cases.

Case 1: Construction of a required map for F_2 .

Let the hyperbolic orbifold S be the \mathbb{RP}^2 with two singular points of indices 2 and 3. A surjective homomorphism $\phi: \pi_1^0(S) = \langle a, c_1, c_2 \mid a^2 c_1 c_2 = c_1^3 = c_2^2 = 1 \rangle \rightarrow Z_{12}$ is defined by $\phi(c_1) = 4$, $\phi(c_2) = 6$, $\phi(a) = 1$ in Z_{12} ; here c_1 and c_2 are presented by orientable loops, a is presented by nonorientable loop. Obviously ϕ is surjective. So ϕ determines a cyclic branched covering $\tilde{S} \rightarrow S$ of degree 12. Since $\phi(c_i)$ and c_i have the same order, kernel ϕ is a surface group by (b). Pick any $\alpha \in \text{kernel } \phi$. Suppose the sums of powers of a, c_1, c_2 in α are k, l, m respectively. Now we must have $k + 4l + 6m = 12j$ for some integer j , so k is even. Thus α is an orientable loop. It follows that \tilde{S} is an orientable surface. From $\chi(\tilde{S}) = -12 \times (1 - \frac{1}{2} - \frac{1}{3}) = -2$, we know $\tilde{S} = F_2$ and the generator s of the deck group of covering is a map of order 12.

Since S has only two singular points of indices 2 and 3, we know that the periodic map s on \tilde{S} has only two periodic orbits of length $\frac{12}{2} = 6$ and $\frac{12}{3} = 4$. So s, s^2, s^3 are all fixed point free.

Case 2: Construction of a required map on F_g , $g > 2$, g is even.

Let the hyperbolic surface S be the connected sum of three \mathbb{RP}^2 . A surjective homomorphism $\phi: \pi_1^0(S) = \pi_1(S) = \langle a, b, c \mid aba^{-1}b^{-1}c^2 = 1 \rangle \rightarrow Z_{2g-2}$ is defined by $\phi(a) = 2$, $\phi(b) = 2$, $\phi(c) = g-1$ in Z_{2g-2} , here a and b are presented by orientable loops and c is presented by a nonorientable loop. Since $g-1$ is odd, the largest common divisor $(g-1, 2) = 1$ and ϕ is surjective. Obviously, kernel ϕ is a surface group and ϕ determines a covering $\tilde{S} \rightarrow S$ of degree $2g-2$ without branched point.

Pick any $\alpha \in \text{kernel } \phi$. Suppose the sum of powers of a, b, c in α are k, l, m respectively. Now we must have $2k + 2l + m(g - 1) = j(2g - 2)$ for some integer j . Since $g - 1$ is odd, m is even, α is an orientable loop. It follows that \tilde{S} is an orientable surface. From $\tilde{\chi}(\tilde{S}) = \chi(\tilde{S}) = (2g - 2) \times \chi(S) = 2 - 2g$, we know $\tilde{S} = F_g$ and the generator s_1 of the deck group of the covering is a map of order $2g - 2$ with free degree $2g - 2$.

Case 3: Construction of a required map for F_g , $g > 2$, g is odd.

Let the hyperbolic orbifold S be the Klein bottle with one singular point of index 2. A surjective homomorphism $\phi: \pi_1^0(S) = \langle a_1, a_2, c \mid a_1^2 a_2^2 c = c^2 = 1 \rangle \rightarrow Z_{4g-4}$ is defined by $\phi(c) = 2g - 2$, $\phi(a_1) = -1$, $\phi(a_2) = g$ in Z_{4g-4} , here a_1, a_2 are presented by nonorientable loops, c is presented by an orientable loop. Obviously ϕ is surjective. So ϕ determines a cyclic branched covering $\tilde{S} \rightarrow S$ of degree $4g - 4$. Since $\phi(c)$ and c have the same order, kernel ϕ is a surface group by (b). Pick any $\alpha \in \text{kernel } \phi$. Suppose the sums of powers of a_1, a_2, c in α are k, l, m respectively. Now we must have $-k + lg + m(2g - 2) = j(4g - 4)$ for some integer j , i.e., $-k + l = j(4g - 4) - m(2g - 2) - l(g - 1)$. Since g is odd, $k + l$ is even. Thus α is an orientable loop. It follows that \tilde{S} is an orientable surface. From $\chi(\tilde{S}) = (4g - 4) \times \chi(S) = (4g - 4) \times (-\frac{1}{2}) = 2 - 2g$, we know $\tilde{S} = F_g$ and the generator s of the deck group of branched covering is a map of order $4g - 4$.

Since S has only one singular point of index 2, we know that the periodic map s on \tilde{S} has only one periodic orbit of length $(4g - 4)/2 = 2g - 2$. So s, s^2, \dots, s^{2g-3} are all fixed point free.

Case 4: Construction of an example on N_3 .

Let a, b be a basis for the first homology group of torus T . There is an orientation preserving periodic map ι' of order 6 determined by the map $(a, b) \mapsto (-b, a + b)$. Now $L(\iota') = 1$, by the elementary fact in fixed point theory, ι' has only one fixed point x . Let D be an invariant small disk of ι' centered at x . The restriction of ι' on $S_1 = T - \text{int } D$ is a map ι of the same order which is fixed point free. Now parameterize ∂S_1 by an angle $0 \leq \theta \leq 2\pi$ such that the restriction of ι on ∂S_1 is a rotation of the angle $2\pi/6$.

Now we identify the antipodal points on ∂S_1 , we get N_3 and ι induces a fixed point free map of order 6 on N_3 .

Case 5: Construction of an example on N_q , $q > 3$.

Pick an oriented essential simple closed curve c on torus T . Let s_1 be a rotation of torus T with angle $2\pi/(q - 2)$ along this curve c and D be a small disk such that $D, s_1(D), \dots, s_1^{q-3}(D)$ are disjoint. Let $S_1 = T - \bigcup_{i=1}^{q-2} s_1^i(D)$. Identifying the antipodal points of each component of ∂S_1 , we get N_q and s_1 induces a periodic map s of order $p - 2$ such that s, s^2, \dots, s^{p-3} are all fixed point free.

Remark. (3) The example in Case 4 cannot be produced by using cyclic branched covering, since s^3 has a one-dimensional fixed point set.

Step 2. To show the numbers in Theorem 1 are the upper bounds of the free degrees on closed surfaces.

The fact that $h(N_q)'s$ and $h_0(F_g)'s$ are bounded by the numbers in Theorem 1 has been proved in [8] and [5] respectively. However we would like to prove the general case rather than to prove the case of orientation reversing homeomorphisms on orientable surfaces only for two reasons: (1) [8] and [5] are not commonly available or not easy to read; (2) the proof for the general case is only few lines longer than the proof of the special case.

Let f be a homeomorphism on a closed surface F with $\chi(F) < 0$ such that f, f^2, \dots, f^n are all fixed point free.

Then we have

$$L(f^{-2}) = L(f^{-1}) = L(f) = L(f^2) = \dots = L(f^n) = 0. \quad (1)$$

Let $\lambda_1, \lambda_2, \dots, \lambda_s$ be the eigenvalues of an induced isomorphism f_{*1} , here s is the rank of $H_1(F, Q)$, then the characteristic polynomial of f_{*1} is

$$P(x) = (x - \lambda_1) \cdots (x - \lambda_s) = x^s - p_1 x^{s-1} + \cdots + (-1)^s p_s, \quad (2)$$

here

$$p_s = \lambda_1 \lambda_2 \cdots \lambda_s = \pm 1. \quad (3)$$

For any nonzero integer, set $\iota_k = \sum_{i=1}^s \lambda_i^k$. Then

$$p_{s-1} = \sum_{i=1}^s \lambda_1 \cdots \hat{\lambda}_i \cdots \lambda_s = \sum_{i=1}^s \frac{1}{\lambda_i} = \iota_{-1}, \quad (4)$$

here $\hat{\lambda}_i$ is the omission of λ_i , and

$$p_{s-2} = \sum_{1 \leq i < j \leq s} \lambda_1 \cdots \hat{\lambda}_i \cdots \hat{\lambda}_j \cdots \lambda_s = \sum_{1 \leq i < j \leq s} \frac{1}{\lambda_i \lambda_j},$$

we have

$$\iota_{-2} = p_{s-1}^2 - 2p_{s-2}. \quad (5)$$

By Newton's formulas about elementary symmetric polynomials (see [7, Exercise 2, Section 29], for example), we have

$$\iota_1 - p_1 = 0, \quad (6.1)$$

$$\iota_2 - p_1 \iota_1 + 2p_2 = 0, \quad (6.2)$$

$$\iota_3 - p_1 \iota_2 + p_2 \iota_1 - 3p_3 = 0, \quad (6.3)$$

and in general

$$\iota_k - p_1 \iota_{k-1} + \cdots + (-1)^{k-1} p_{k-1} \iota_1 + (-1)^k k p_k = 0, \quad 1 \leq k \leq s. \quad (6.k)$$

With all those elementary facts in hand, we are going to prove the second step.

Case 1: Suppose f is a homeomorphism on F_g such that f, \dots, f^n are all fixed point free. We show $n < 2g$ and $n < 2g - 2$ if further $g > 2$. Now $s = 2g$.

If f is orientation reversing, then f^{2k-1} is a self map of degree -1 and f^{2k} is a self map of degree 1 for any integer k . From (1), linear algebra and the definition of the Lefschetz number, we have

$$\iota_{2k} = \sum_{i=1}^{2g} \lambda_i^{2k} = 2, \quad -2 \leq 2k \leq n, \quad k \neq 0, \quad (7.2k)$$

$$\iota_{2k-1} = \sum_{i=1}^{2g} \lambda_i^{2k-1} = 0, \quad 0 \leq 2k \leq n. \quad (7.2k-1)$$

By (7.i), (6.1) is $p_1 = 0$, and then (6.2) is $p_2 = -1$. We use induction to show $p_l = 0, 3 \leq l \leq s$. Now (6.3) is $p_3 = 0$. Suppose $p_k = 0, 3 < k < n$, then (6.k+1) has the form

$$\iota_{k+1} + (-1)\iota_{k-1} + (k+1)p_{k+1} = 0.$$

If k is even, then we have $(k+1)p_{k+1} = 0$, so $p_{k+1} = 0$. If k is odd, then we have $2 - 2 + (k+1)p_{k+1} = 0$, so $p_{k+1} = 0$.

It f is orientation preserving, then f^k is a self map of degree 1 for any integer k . Similarly we have

$$\iota_k = \sum_{i=1}^{2g} \lambda_i^k = 2, \quad -2 \leq k \leq n, \quad k \neq 0. \quad (8.k)$$

By (8.k), (6.1) is $p_1 = 2$, and then (6.2) is $p_2 = 1$, and then $p_3 = p_4 = \dots = p_s = 0$ by (6.k), $k = 3, \dots, s$.

If $n \geq 2g$, since $2g \geq 3$, $p_{2g} = 0$. This contradicts (3).

If $g > 2$ and $n \geq 2g - 2$, since $2g - 2 \geq 3$, we have $p_{2g-2} = p_{2g-1} = 0$. By (5), we have $\iota_{-2} = 0$, this contradicts (7.-2) and (8.-2).

Case 2: Suppose f is a homeomorphism on N_q such that f, \dots, f^n are all fixed point free. We show $n < q - 1$ and $n < q - 2$ if further $q > 3$. Now $s = q - 1$.

Since $H_2(N_g, Q) = 0$, similarly we have

$$\iota_k = \sum_{i=1}^{2g} \lambda_i^k = 1, \quad -1 \leq k \leq n, \quad k \neq 0. \quad (9.k)$$

By (9.k), (6.1) is $p_1 = 1$, and then $p_2 = p_3 = p_4 = \dots = p_s = 0$ by (6.k), $k = 2, \dots, s$. If $n \geq q - 1$, since $q > 2$, $p_{q-1} = 0$. This contradicts (3).

If $q > 3$ and $n \geq q - 2$, since $q - 2 \geq 2$, we have $p_{q-2} = 0$. By (4), we have $\iota_{-1} = 0$. This contradicts (9.-1).

We have finished the proof of Theorem 1. \square

The author thanks the referee for his comments which improve the presentation of Step 2 significantly.

References

- [1] I. Bernstein and A. Edmonds, On the construction of branch coverings of low-dimensional manifolds, Trans. Amer. Math. Soc. 247 (1979) 87-125.

- [2] R. Brown, *The Lefschetz Fixed Point Theorem* (Scott, Foresman, Glenview, IL, 1971).
- [3] F. Fuller, The existence of periodic points, *Ann. of Math.* 57 (1953) 229–230.
- [4] W.J. Harvey, Cyclic group of automorphisms of compact Riemann surface, *Quart. J. Math. Oxford Ser. (2)* 17 (1966) 86–97.
- [5] J. Nielsen, Fixed point free mappings, *Mat. Tidsskr. B* (1942) 25–41 (in Danish); reviewed by R. Fox, *Math. Reviews* 7 (1946) 137.
- [6] P. Scott, The geometry of 3-manifolds, *Bull. London Math. Soc.* 15 (1983) 401–487.
- [7] Van der Waerden, *Algebra I* (Springer, Berlin, 4th ed., 1955).
- [8] S. Wang, Free degree of homeomorphisms on non-orientable closed surfaces, *Kexue Tongbao (Chinese)* 28 (1983) 1156–1158.
- [9] S. Wang, Maximum orders of periodic maps on closed surfaces, *Topology Appl.* 41 (1991) 255–262.